

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

71

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification: A61K 31/415	A1	(11) International Publication Number: WO 99/61019 (43) International Publication Date: 2 December 1999 (02.12.99)
(21) International Application Number: PCT/US99/11322 (22) International Filing Date: 21 May 1999 (21.05.99) (30) Priority Data: 60/086,494 22 May 1998 (22.05.98) US (71) Applicant: AVANIR PHARMACEUTICALS [US/US]; 9393 Towne Centre Drive #200, San Diego, CA 92121 (US). (72) Inventors: SIRCAR, Jagadish; Avanir Pharmaceuticals, 9393 Towne Centre Drive #200, San Diego, CA 92121 (US). RICHARDS, Mark, L.; Avanir Pharmaceuticals, 9393 Towne Centre Drive #200, San Diego, CA 92121 (US). CAMPBELL, Michael, G.; Avanir Pharmaceuticals, 9393 Towne Centre Drive #200, San Diego, CA 92121 (US). MAJOR, Michael, W.; Avanir Pharmaceuticals, 9393 Towne Centre Drive #200, San Diego, CA 92121 (US). (74) Agent: ALTMAN, Daniel, E.; Knobbe, Martens, Olson & Bear, LLP, 16th floor, 620 Newport Center Drive, Newport Beach, CA 92660 (US).		(81) Designated States: AE, AL, AM, AT, AT (Utility model), AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, CZ (Utility model), DE, DE (Utility model), DK, DK (Utility model), EE, EE (Utility model), ES, FI, FI (Utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (Utility model), SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: BENZIMIDAZOLE DERIVATIVES AS MODULATORS OF IgE		
(57) Abstract This invention relates to a family of diacyl benzimidazole analogs, which are inhibitors of the IgE response to allergens. These compounds are useful in the treatment of allergy and/or asthma or any diseases where IgE is pathogenic.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CJ	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
ER	Estonia						

BENZIMIDAZOLE DERIVATIVES AS MODULATORS OF IgE

Background of the Invention

This invention relates to small molecule inhibitors of the IgE response to allergens that are useful in the treatment of allergy and/or asthma or any diseases where IgE is pathogenic.

An estimated 10 million persons in the United States have asthma, about 5% of the population. The estimated cost of asthma in the United States exceeds \$6 billion. About 25% of patients with asthma who seek emergency care require hospitalization, and the largest single direct medical expenditure for asthma has been inpatient hospital services (emergency care), at a cost of greater than \$1.6 billion. The cost for prescription medications, which increased 54% between 1985 and 1990, was close behind at \$1.1 billion (Kelly, *Pharmacotherapy* 12:13S-21S (1997)).

According to the National Ambulatory Medical Care Survey, asthma accounts for 1% of all ambulatory care visits, and the disease continues to be a significant cause of missed school days in children. Despite improved understanding of the disease process and better drugs, asthma morbidity and mortality continue to rise in this country and worldwide (U.S. Department of Health and Human Services; 1991, publication no. 91-3042). Thus, asthma constitutes a significant public health problem.

The pathophysiologic processes that attend the onset of an asthmatic episode can be broken down into essentially two phases, both marked by bronchoconstriction, that causes wheezing, chest tightness, and dyspnea. The first, early phase asthmatic response is triggered by allergens, irritants, or exercise. Allergens cross-link immunoglobulin E (IgE) molecules bound to receptors on mast cells, causing them to release a number of pre-formed inflammatory mediators, including histamine. Additional triggers include the osmotic changes in airway tissues following exercise or the inhalation of cold, dry air. The second, late phase response that follows is characterized by infiltration of activated eosinophils and other inflammatory cells into airway tissues, epithelial desquamation, and by the presence of highly viscous mucus within the airways. The damage caused by this inflammatory response leaves the airways "primed" or sensitized, such that smaller triggers are required to elicit subsequent asthma symptoms.

A number of drugs are available for the palliative treatment of asthma; however, their efficacies vary markedly. Short-acting β_2 -adrenergic agonists, terbutaline and albuterol, long the mainstay of asthma treatment, act primarily during the early phase as bronchodilators. The newer

long-acting β_2 -agonists, salmeterol and formoterol, may reduce the bronchoconstrictive component of the late response. However, because the β_2 -agonists do not possess significant antiinflammatory activity, they have no effect on bronchial hyperreactivity.

Numerous other drugs target specific aspects of the early or late asthmatic responses. For example, antihistamines, like loratadine, inhibit early histamine-mediated inflammatory responses. Some of the newer antihistamines, such as azelastine and ketotifen, may have both antiinflammatory and weak bronchodilatory effects, but they currently do not have any established efficacy in asthma treatment. Phosphodiesterase inhibitors, like theophylline/xanthines, may attenuate late inflammatory responses, but there is no evidence that these compounds decrease bronchial hyperreactivity. Anticholinergics, like ipratropium bromide, which are used in cases of acute asthma to inhibit severe bronchoconstriction, have no effect on early or late phase inflammation, no effect on bronchial hyperreactivity, and therefore, essentially no role in chronic therapy.

The corticosteroid drugs, like budesonide, are the most potent antiinflammatory agents. Inflammatory mediator release inhibitors, like cromolyn and nedocromil, act by stabilizing mast cells and thereby inhibiting the late phase inflammatory response to allergen. Thus, cromolyn and nedocromil, as well as the corticosteroids, all reduce bronchial hyperreactivity by minimizing the sensitizing effect of inflammatory damage to the airways. Unfortunately, these antiinflammatory agents do not produce bronchodilation.


Several new agents are currently being developed that inhibit specific aspects of asthmatic inflammation. For instance, leukotriene receptor antagonists (ICI-204, 219, accolate), specifically inhibit leukotriene-mediated actions. The leukotrienes have been implicated in the production of both airway inflammation and bronchoconstriction.

Thus, while numerous drugs are currently available for the treatment of asthma, these compounds are primarily palliative and/or have significant side effects. Consequently, new therapeutic approaches which target the underlying cause rather than the cascade of symptoms would be highly desirable. Asthma and allergy share a common dependence on IgE-mediated events. Indeed, it is known that excess IgE production is the underlying cause of allergies in general and allergic asthma in particular (Duplantier and Cheng, *Ann. Rep. Med. Chem.* 29:73-81 (1994)). Thus, compounds that lower IgE levels may be effective in treating the underlying cause of asthma and allergy.

None of the current therapies eliminate the excess circulating IgE. The hypothesis that lowering plasma IgE may reduce the allergic response, was confirmed by recent clinical results with chimeric anti-IgE antibody, CGP-51901, and recombinant humanized monoclonal antibody, rhuMAB-E25. Indeed, three companies, Tanox Biosystems, Inc., Genentech Inc. and Novartis AG are collaborating in the development of a humanized anti-IgE antibody (BioWorld® Today, February 26, 1997, p. 2) which will treat allergy and asthma by neutralizing excess IgE. Tanox has already successfully tested the anti-IgE antibody, CGP-51901, which reduced the severity and duration of nasal symptoms of allergic rhinitis in a 155-patient Phase II trial (Scrip #2080, Nov 24, 1995, p.26). Genentech recently disclosed positive results from a 536 patient phase-II/III trials of its recombinant humanized monoclonal antibody, rhuMAB-E25 (BioWorld® Today, November 10, 1998, p. 1). The antibody, rhuMAB-E25, administered by injection (highest dose 300 mg every 2 to 4 weeks as needed) provided a 50% reduction in the number of days a patient required additional "rescue" medicines (antihistamines and decongestants), compared to placebo. An NDA filing for this product is projected to be in the year 2000. The positive results from anti-IgE antibody trials suggest that therapeutic strategies aimed at IgE down-regulation may be effective.

Summary of the Invention

The present invention discloses a family of related compounds for use in the treatment of a condition associated with an excess IgE level. The benzimidazole inhibitors of IgE in accordance with the present invention are represented by the generic formula:



X and Y are independently selected from the group consisting of H, alkyl, alkoxy, aryl, substituted aryl, hydroxy, halogen, amino, alkylamino, nitro, cyano, CF₃, OCF₃, CONH₂, CONHR and NHCOR₁. R is selected from the group consisting of H, CH₃, C₂H₅, C₃H₇, C₄H₉,

CH_2Ph , and $\text{CH}_2\text{C}_6\text{H}_4\text{-F(p-)}$. R_1 and R_2 are independently selected from the group consisting of H, aryl, substituted aryl, cycloaryl substituted cycloaryl, multi-ring cycloaryl, benzyl, substituted benzyl and the like. Substitutions are alkyl, aryl, CF_3 , CH_3 , OCH_3 , OH, CN, COOR, COOH and the like.

In accordance with another aspect of the invention, there is disclosed a composition for use in the treatment of an allergic condition comprising the diacyl benzimidazole inhibitor of IgE disclosed above and at least one additional active ingredient, combined in a pharmaceutically acceptable diluent. The additional active ingredients may be selected from the group consisting of short-acting β_2 -adrenergic agonists, like terbutaline and albuterol, long-acting β_2 -adrenergic agonists, like salmeterol and formoterol, antihistamines, like loratadine, azelastine and ketotifen, phosphodiesterase inhibitors, anticholinergic agents, corticosteroids, inflammatory mediator release inhibitors and leukotriene receptor antagonists.

In accordance with another aspect of the invention, there is disclosed a family of symmetric and asymmetric diacyl and monoacyl benzimidazole compounds for use in the treatment of an allergic condition comprising the following species:

Structure 	BIAA
	$C_{27}H_{20}N_4O_2$
	CLOGP 5.47

Structure 	BIAB
	$C_{27}H_{19}ClN_4O_2$
	CLOGP 6.24

Structure 	BIAC
	$C_{28}H_{22}N_4O_3$
	CLOGP 5.58

Structure 	BIAD
	$C_{26}H_{19}N_5O_2$
	CLOGP 4.28

Structure 	BIAE
	$C_{30}H_{26}N_4O_5$
	CLOGP 4.82

Structure 	BIAF
	$C_{27}H_{18}Cl_2N_4O_2$
	CLOGP 6.97

Structure 	BIAG
	$C_{27}H_{18}Cl_2N_4O_2$
	CLOGP 5.31













Structure 	BIAH
	$C_{27}H_{18}Cl_2N_4O_2$
	CLOGP 6.14













Structure 	BIAI
	$C_{27}H_{26}N_4O_2$
	CLOGP 6.01


Structure 	BIAJ
	$C_{26}H_{24}N_4O_2$
	CLOGP 5.45


Structure 	BIAK
	$C_{25}H_{18}N_4O_2S$
	CLOGP 5.20


Structure 	BIAL
	$C_{25}H_{18}N_4O_2S$
	CLOGP 5.20








Structure	BICA
	$C_{28}H_{22}N_4O_3$
	CLOGP
	5.58


Structure	BICB
	$C_{28}H_{21}ClN_4O_3$
	CLOGP
	6.35


Structure 	BICC
	$C_{29}H_{24}N_4O_4$
	CLOGP 5.70


Structure 	BICD
	$C_{27}H_{21}N_5O_3$
	CLOGP 4.39


Structure	BICE
	$C_{31}H_{28}N_4O_6$
	CLOGP
	4.93


Structure	BICF
	$C_{28}H_{20}Cl_2N_4O_3$
	CLOGP
	7.09


Structure	BICG
	$C_{28}H_{20}Cl_2N_4O_3$
	CLOGP
	5.43













Structure 	BICH
	$C_{28}H_{20}Cl_2N_4O_3$
	CLOGP
	6.26













Structure	BICI
	$C_{28}H_{28}N_4O_3$
	CLOGP
	6.12













Structure	BICJ
	$C_{27}H_{26}N_4O_3$
	CLOGP
	5.56













Structure	BICK
	$C_{26}H_{20}N_4O_3S$
	CLOGP
	5.32













Structure	BICL
	$C_{26}H_{20}N_4O_3S$
	CLOGP
	5.32











Structure 	BIGA
	$C_{27}H_{18}Cl_2N_4O_2$
	CLOGP
	5.34

Structure 	BIGB
	$C_{27}H_{17}Cl_3N_4O_2$
	CLOGP
	6.12

Structure 	BIGC
	$C_{28}H_{20}Cl_2N_4O_3$
	CLOGP
	5.46

Structure 	BIGD
	$C_{26}H_{17}Cl_2N_5O_2$
	CLOGP
	4.16

Structure 	BIGE
	$C_{30}H_{24}Cl_2N_4O_5$
	CLOGP
	4.69

Structure 	BIGF
	$C_{27}H_{16}Cl_4N_4O_2$
	CLOGP
	6.85

Structure 	BIGG
	$C_{27}H_{16}Cl_4N_4O_2$
	CLOGP
	5.19













Structure 	BIGH
	$C_{27}H_{16}Cl_4N_4O_2$
	CLOGP
	6.02













Structure 	BIGI
	$C_{27}H_{24}Cl_2N_4O_2$
	CLOGP
	5.88













Structure 	BIGJ
	$C_{26}H_{22}Cl_2N_4O_2$
	CLOGP
	5.33













Structure 	BIGK
	$C_{25}H_{16}Cl_2N_4O_2$
	S
	CLOGP
5.08	













Structure 	BIGL
	$C_{25}H_{16}Cl_2N_4O_2$
	S
	CLOGP
5.08	
















Structure 	BIJG
	$C_{26}H_{22}Cl_2N_4O_2$
	CLOGP 5.29


Structure 	BIJH
	$C_{26}H_{22}Cl_2N_4O_2$
	CLOGP 6.12


avanlr_vlib.db


Structure 	BIIA
	$C_{27}H_{26}N_4O_2$
	CLOGP 6.01


Structure 	BIIB
	$C_{27}H_{25}ClN_4O_2$
	CLOGP 6.78


Structure 	BIIC
	$C_{28}H_{28}N_4O_3$
	CLOGP 6.12


Structure 	BIID
	$C_{26}H_{25}N_5O_2$
	CLOGP 4.82


Structure 	BIIIE
	$C_{30}H_{32}N_4O_5$
	CLOGP 5.36


Structure 	BIIF
	$C_{27}H_{24}Cl_2N_4O_2$
	CLOGP 7.51


Structure 	BIIG
	$C_{27}H_{24}Cl_2N_4O_2$
	CLOGP 5.85


Structure 	BIIH
	$C_{27}H_{24}Cl_2N_4O_2$
	CLOGP 6.68


Structure 	BIJK
	$C_{25}H_{24}N_4O_2S$
	CLOGP 6.74


Structure 	BIIL
	$C_{25}H_{24}N_4O_2S$
	CLOGP 5.74


Structure 	BIJA
	$C_{26}H_{24}N_4O_2$
	CLOGP 5.45


Structure 	BIJB
	$C_{26}H_{23}ClN_4O_2$
	CLOGP 6.22


Structure 	BIJC
	$C_{27}H_{26}N_4O_3$
	CLOGP 5.56

Structure 	BIJD
	$C_{25}H_{23}N_5O_2$
	CLOGP 4.26


Structure 	BIJE
	$C_{29}H_{30}N_4O_5$
	CLOGP 4.80


Structure 	BIJF
	$C_{26}H_{22}Cl_2N_4O_2$
	CLOGP 6.95


Structure 	BIOK
	$C_{29}H_{28}N_4O_2S$
	CLOGP
	6.37


Structure 	BIOL
	$C_{29}H_{28}N_4O_2S$
	CLOGP
	6.37


avanir_vlib.db


Structure 	BIOA
	$C_{31}H_{30}N_4O_2$
	CLOGP
	6.64


Structure 	BIOB
	$C_{31}H_{29}ClN_4O_2$
	CLOGP
	7.41


Structure 	BIOG
	$C_{32}H_{32}N_4O_3$
	CLOGP
	6.75











Structure 	BIOD
	$C_{30}H_{29}N_4O_2$
	CLOGP
	5.45











Structure 	BIOE
	$C_{34}H_{36}N_4O_5$
	CLOGP
	5.99











Structure 	BIOF
	$C_{31}H_{28}Cl_2N_4O_2$
	CLOGP
	8.14

Structure 	BIOG
	$C_{31}H_{28}Cl_2N_4O_2$
	CLOGP
	6.48


Structure 	BIOH
	$C_{31}H_{28}Cl_2N_4O_2$
	CLOGP
	7.31








In accordance with another aspect of the present invention, there is disclosed a method for the preparation of a medicament for treatment of a condition associated with an excess IgE level. The compound has the formula:



X and Y are independently selected from the group consisting of H, alkyl, alkoxy, aryl, substituted aryl, hydroxy, halogen, amino, alkylamino, nitro, cyano, CF₃, OCF₃, CONH₂, CONHR and NHCOR₁. R is selected from the group consisting of H, CH₃, C₂H₅, C₃H₇, C₄H₉, CH₂Ph, and CH₂C₆H₄-F(p-). R₁ and R₂ are independently selected from the group consisting of H, aryl, substituted aryl, cycloaryl substituted cycloaryl, multi-ring cycloaryl, benzyl, substituted benzyl and the like. Substitutions are alkyl, aryl, CF₃, CH₃, OCH₃, OH, CN, COOR, COOH and the like.

In accordance with another aspect of the present invention, there is disclosed a method of treating a mammal having a condition associated with an excess IgE level. The method comprises administering to the mammal an amount of a compound sufficient to reduced IgE levels in the mammal. The compound has the formula:



X and Y are independently selected from the group consisting of H, alkyl, alkoxy, aryl, substituted aryl, hydroxy, halogen, amino, alkylamino, nitro, cyano, CF₃, OCF₃, CONH₂, CONHR and NHCOR₁. R is selected from the group consisting of H, CH₃, C₂H₅, C₃H₇, C₄H₉,

CH₂Ph, and CH₂C₆H₄-F(p-). R₁ and R₂ are independently selected from the group consisting of H, aryl, substituted aryl, cycloaryl substituted cycloaryl, multi-ring cycloaryl, benzyl, substituted benzyl, alkyl, cycloalkyl substituted cycloalkyl, multi-ring cycloalkyl, fused-ring aliphatic, cyclopropyl, substituted cyclopropyl, cyclobutyl, substituted cyclobutyl, cyclopentyl, substituted cyclopentyl, cyclohexyl, substituted cyclohexyl, cycloheptyl, substituted cycloheptyl, bicycloheptyl, bicyclooctyl, bicyclononyl, substituted bicycloalknyl, adamantyl, substituted adamantyl and the like, wherein at least one of R₁ and R₂ are aromatic groups. Substitutions are alkyl, aryl, CF₃, CH₃, OCH₃, OH, CN, COOR, COOH and the like.

In a variation of the above-disclosed method, at least one additional active ingredient may be administered in conjunction with the administration of the compound. The additional active ingredient may be combined with said compound in a pharmaceutically acceptable diluent and co-administered to the mammal. The additional active ingredient may be a short-acting β_2 -adrenergic agonist selected from the group consisting of terbutaline and albuterol. In a variation, the additional active ingredient may be a long-acting β_2 -adrenergic agonist selected from the group consisting of salmeterol and formoterol or an antihistamine selected from the group consisting of loratadine, azelastine and ketotifen. In another variation, the additional active ingredient may be a phosphodiesterase inhibitor, an anticholinergic agent, a corticosteroid, an inflammatory mediator release inhibitor or a leukotriene receptor antagonist.

The compound is preferably administered at a dose of about 0.01 mg to about 100 mg per kg body weight per day in divided doses of said compound for at least two consecutive days at regular periodic intervals.

Other variations within the scope of the present invention may be more fully understood with reference to the following detailed description.

Detailed Description of the Preferred Embodiment

The present invention is directed to small molecule inhibitors of IgE (synthesis and/or release) which are useful in the treatment of allergy and/or asthma or any diseases where IgE is pathogenic. The particular compounds disclosed herein were identified by their ability to suppress IgE levels in both *ex vivo* and *in vivo* assays. Development and optimization of clinical treatment regimens can be monitored by those of skill in the art by reference to the *ex vivo* and *in vivo* assays described below.

Ex Vivo Assay

This assay begins with *in vivo* antigen priming and measures secondary antibody responses *in vitro*. The basic protocol was documented and optimized for a range of parameters including: antigen dose for priming and time span following priming, number of cells cultured *in vitro*, antigen concentrations for eliciting secondary IgE (and other Ig's) response *in vitro*, fetal bovine serum (FBS) batch that will permit optimal IgE response *in vitro*, the importance of primed CD4+ T cells and hapten-specific B cells, and specificity of the ELISA assay for IgE (Marcelletti and Katz, *Cellular Immunology* 135:471-489 (1991); incorporated herein by reference).

The actual protocol utilized for this project was adapted for a more high throughput analyses. BALB/cByj mice were immunized i.p. with 10 µg DNP-KLH adsorbed onto 4 mg alum and sacrificed after 15 days. Spleens were excised and homogenized in a tissue grinder, washed twice, and maintained in DMEM supplemented with 10% FBS, 100 U/ml penicillin, 100 µg/ml streptomycin and 0.0005% 2-mercaptoethanol. Spleen cell cultures were established (2-3 million cells/ml, 0.2 ml/well in quadruplicate, 96-well plates) in the presence or absence of DNP-KLH (10 ng/ml). Test compounds (2 µg/ml and 50 ng/ml) were added to the spleen cell cultures containing antigen and incubated at 37° C for 8 days in an atmosphere of 10% CO₂.

Culture supernatants were collected after 8 days and Ig's were measured by a modification of the specific isotype-selective ELISA assay described by Marcelletti and Katz (*Supra*). The assay was modified to facilitate high throughput. ELISA plates were prepared by coating with DNP-KLH overnight. After blocking with bovine serum albumin (BSA), an aliquot of each culture supernatant was diluted (1:4 in phosphate buffered saline (PBS) with BSA, sodium azide and Tween 20), added to the ELISA plates, and incubated overnight in a humidified box at 4° C. IgE levels were quantitated following successive incubations with biotinylated-goat antimouse IgE (b-GAME), AP-streptavidin and substrate.

Antigen-specific IgG1 was measured similarly, except that culture supernatants were diluted 200-fold and biotinylated-goat antimouse IgG1 (b-GAMG1) was substituted for b-GAME. IgG2a was measured in ELISA plates that were coated with DNP-KLH following a 1:20 dilution of culture supernatants and incubation with biotinylated-goat antimouse IgG2a (b-GAMG2a). Quantitation of each isotype was determined by comparison to a standard curve. The level of detectability of all

antibody was about 200-400 pg/ml and there was less than 0.001% cross-reactivity with any other Ig isotype in the ELISA for IgE.

In Vivo Assay


Compounds found to be active in the *ex vivo* assay (above) were further tested for their activity in suppressing IgE responses *in vivo*. Mice receiving low-dose radiation prior to immunization with a carrier exhibited an enhanced IgE response to sensitization with antigen 7 days later. Administration of the test compounds immediately prior to and after antigen sensitization, measured the ability of that drug to suppress the IgE response. The levels of IgE, IgG1 and IgG2a in serum were compared.

Female BALB/cByj mice were irradiated with 250 rads 7 hours after initiation of the daily light cycle. Two hours later, the mice were immunized i.p. with 2 μ g of KLH in 4 mg alum. Two to seven consecutive days of drug injections were initiated 6 days later on either a once or twice daily basis. Typically, i.p. injections and oral gavages were administered as suspensions (150 μ l/injection) in saline with 10% ethanol and 0.25% methylcellulose. Each treatment group was composed of 5-6 mice. On the second day of drug administration, 2 μ g of DNP-KLH was administered i.p. in 4 mg alum, immediately following the morning injection of drug. Mice were bled 7-21 days following DNP-KLH challenge.

Antigen-specific IgE, IgG1 and IgG2a antibodies were measured by ELISA. Periorbital bleeds were centrifuged at 14,000 rpm for 10 min, the supernatants were diluted 5-fold in saline, and centrifuged again. Antibody concentrations of each bleed were determined by ELISA of four dilutions (in triplicate) and compared to a standard curve: anti-DNP IgE (1:100 to 1:800), anti-DNP IgG2a (1:100 to 1:800), and anti-DNP IgG1 (1:1600 to 1:12800).

Diacyl Benzimidazole Inhibitors of IgE


Several species embraced by the following generic formula were synthesized and evaluated for their effectiveness in down-regulating IgE in the *ex vivo* and *in vivo* assays.



X and Y are independently selected from the group consisting of H, alkyl, alkoxy, aryl, substituted aryl, hydroxy, halogen, amino, alkylamino, nitro, cyano, CF_3 , OCF_3 , CONH_2 , CONHR and NHCOR_1 . R is selected from the group consisting of H, CH_3 , C_2H_5 , C_3H_7 , C_4H_9 , CH_2Ph , and $\text{CH}_2\text{C}_6\text{H}_4\text{-F(p-)}$. R_1 and R_2 are independently selected from the group consisting of H, aryl, substituted aryl, cycloaryl substituted cycloaryl, multi-ring cycloaryl, benzyl, substituted benzyl, alkyl, cycloalkyl substituted cycloalkyl, multi-ring cycloalkyl, fused-ring aliphatic, cyclopropyl, substituted cyclopropyl, cyclobutyl, substituted cyclobutyl, cyclopentyl, substituted cyclopentyl, cyclohexyl, substituted cyclohexyl, cycloheptyl, substituted cycloheptyl, bicycloheptyl, bicyclooctyl, bicyclononyl, substituted bicycloalknyl, adamantyl, substituted adamantyl and the like, wherein at least one of R_1 and R_2 are aromatic groups. Substitutions are alkyl, aryl, CF_3 , CH_3 , OCH_3 , OH , CN , COOR , COOH and the like.

Synthesis of the Combinatorial Library

The diacyl benzimidazole compounds of the present invention were prepared using the following synthesis reactions, wherein the desired acid chlorides are selected from the R_1 and R_2 groups provided in the Table.



Synthesis of 3: 4-Nitro-1,2-phenylenediamine (10 g, 65.3 mmol) and 4-aminobenzoic acid (8.95 g, 65.3 mmol) were taken in a round bottomed flask and phosphorus oxychloride (95 ml) was added slowly. The reaction mixture was allowed to stir under reflux conditions. After 18 h, the reaction was allowed to cool and then poured slowly into an ice water mixture in an Erlenmeyer flask with vigorous stirring. Greenish yellow precipitate fell out which was then

filtered and washed with copious amounts of water. The residue was then dried to obtain 16.9 g of crude desired product. Mass spectrum analysis (positive ion) indicated presence of 3.

Synthesis of 4: Benzimidazole 3 (800 mg, 3.14 mmol) was dissolved in dry pyridine (5 ml) in a scintillation vial and the desired acid chlorides (1.1 eq) were added slowly. The reactions were carried out in an oven at 60°C. After 16h, the reaction was cooled to RT and DI water was added. Precipitation took place, which was filtered off, washed with water and air dried. The aqueous layer was extracted with EtOAc (6 x 50 ml), dried over anhydrous Na₂SO₄ and the solvent was removed *in vacuo* to result in a colored solid. By positive ion MS the desired monoacylated product was found to be present in the initial precipitate as well as in the organic layer. Hence the solid residues obtained were combined and used as such for the reduction step.


Reduction of 4: Crude monoacylated nitro benzimidazole 4 (1.22 g, 3.40 mmol) was dissolved in MeOH (20 ml) and minimum amount of THF was added for complete dissolution to occur. Catalytic amount of 10% Pd on C was added and the solution was degassed and allowed to stir at 3.4 atm pressure under H₂ atmosphere for 4 h. Upon completion of reaction as observed via TLC, the reaction mixture was filtered through celite and the solvent was removed under reduced pressure to afford 979 mg of crude residue.

General Organic Analyses

HPLC/MS data was obtained using a Gilson semi-prep HPLC with a Gilson 170 Diode Array UV detector and PE Sciex API 100LC MS based detector. A Waters 600E with a Waters 490E UV detector was also used for recording HPLC data. The compounds were eluted with a gradient of CH₃CN (with 0.0035% TFA) and H₂O (with 0.01% TFA). Both HPLC instruments used Advantage C18 60A 5μ 50mm x 4.6mm columns from Thomson Instrument Company. Mass spectra were obtained by direct injection and electrospray ionization on a PE Sciex API 100LC MS based detector. Thin layer chromatography was performed using Merck 60F-254 aluminum backed precoated plates. Flash chromatography was carried out on Merck silica gel 60 (230-400 mesh) purchased from EM Scientific.

Syntheses of Symmetrical Diamides


The symmetrical diacyl benzimidazole compounds of the present invention were generally prepared from 2-(4-aminophenyl)-5-aminobenzimidazole, which was obtained by reduction of 2-(4-nitrophenyl)-6-nitrobenzimidazole.



2-(4-nitrophenyl)-6-nitrobenzimidazole

The dinitro benzimidazole was prepared as follows: a mixture of 4-nitrophenylenediamine (6.4g, 41.83 mmol) and 4-nitrobenzoic acid (7.86 g, 47 mmol) was dissolved in POCl_3 (250 ml) and heated to reflux for 2 h. The reaction mixture was cooled, poured on to ice, and stirred for 30 min. The resulting solid was filtered and washed with methanol and sodium bicarbonate to remove unreacted acid and allowed to dry overnight to give the desired product as a brown solid (5.8 g). The product was characterized by electrospray mass spectroscopy (mp $>300^\circ\text{C}$).

2-(4-Aminophenyl)-5-aminobenzimidazole was prepared by suspending the above solid (75 g) in THF (75 ml), to which was added Pd-C (10% Pd by weight). The flask was purged with hydrogen and stirred under a balloon of hydrogen over night. TLC and MS showed starting material was still present so the reaction was allowed to continue over the weekend. TLC indicated complete reaction, the reaction was filtered through celite and washed with methanol. The solvent was removed under reduced pressure to give a dark brown solid (0.37 g) that was used without further purification.




2-(4-aminophenyl)-5-aminobenzimidazole

Alternatively, the 2-(4-aminophenyl)-5-aminobenzimidazole was prepared by the following reduction: 2-(4-nitrophenyl)-6-nitrobenzimidazole (8.9 g, 31 mmole) was suspended in concentrated HCl (100 ml) to which was added stannous chloride (42.3 g 180 mmole). The reaction mixture was heated to reflux for 5 hrs. The mixture was cooled to RT and the HCl salt


of the desired product was precipitated by the addition of ethanol. The resulting solid was filtered, re-dissolved in water and the solution made basic by the addition of concentrated ammonium hydroxide. The resulting precipitate was filtered and dried overnight under vacuum to yield the desired product as a gray solid (6.023 g, 26.9 mmole, 87%). The product characterized by electrospray mass spectroscopy and HPLC (mp. 222-227° C).

2-(4-Aminophenyl)-5-methoxy benzimidazole was synthesized from 2-(4-nitrophenyl)-5-methoxy benzimidazole, which was prepared as follows: 1,2-diamino-4-methoxybenzene (1.26 g, 10.0 mmole) was mixed with 4-nitrobenzoic acid (1.67 g, 9.8 mmole) and dissolved in POCl₃ (10 ml) and heated to reflux for 2.5 hours. The reaction mixture was cooled and cautiously poured onto ice. The resulting solid was filtered, washed with NaHCO₃ and used without further purification.



2-(4-nitrophenyl)-5-methoxy benzimidazole


2-(4-Aminophenyl)-5-methoxy benzimidazole was prepared by dissolving 1 g of the above nitrobenzimidazole in 30% Na₂S•9H₂O (20 ml) with stirring at RT for 21 h. The reaction mixture was diluted with water and extracted with EtOAc. The combined organic extracts were dried over sodium sulfate and concentrated under vacuum. The product was characterized by mass spectroscopy.



2-(4-aminophenyl)-5-methoxy benzimidazole


2-(4-Aminophenyl)-5,6-dichloro benzimidazole was synthesized from 2-(4-nitrophenyl)-5,6-dichloro benzimidazole, which was prepared as follows: 1,2-diamino-4,5-dichlorobenzene (1.68 g, 10.0 mmole) was mixed with 4-nitrobenzoic acid (1.58 g, 9.3 mmole), dissolved in POCl₃ (10 ml), and heated to reflux for 2.5 hours. The reaction mixture was cooled and

cautiously poured onto ice. The resulting solid was filtered, washed with NaHCO_3 and used without further purification.




2-(4-nitrophenyl)-5,6-dichloro benzimidazole

2-(4-Aminophenyl)-5,6-dichloro benzimidazole was prepared by dissolving 1 g of the above nitrobenzimidazole in 30% $\text{Na}_2\text{S}\cdot 9\text{H}_2\text{O}$ (20 ml) with stirring at RT for 21 h. The reaction mixture was diluted with water and extracted with EtOAc. The combined organic extracts were dried over sodium sulfate and concentrated under vacuum. The product was characterized by mass spectroscopy.



2-(4-Aminophenyl)-5,6-dichloro benzimidazole


2-(4-aminophenyl)-7-methyl benzimidazole was synthesized from 2-(4-nitrophenyl)-7-methyl benzimidazole, which was prepared by mixing 1,2-diamino-3-methylbenzene (1.24 g, 10.0 mmole) with 4-nitrobenzoic acid (1.69 g, 9.8 mmole), dissolved in POCl_3 (10 ml), and heated to reflux for 2.5 hours. The reaction mixture was cooled and cautiously poured onto ice. The resulting solid was filtered, washed with NaHCO_3 and used without further purification.



2-(4-nitrophenyl)-7-methyl benzimidazole


2-(4-Aminophenyl)-7-methyl benzimidazole was synthesized by dissolving 1 g of the above nitrobenzimidazole in 30% $\text{Na}_2\text{S}\cdot 9\text{H}_2\text{O}$ (20 ml) with stirring at RT for 4.5 h. The reaction mixture was diluted with water and extracted with EtOAc. The combined organic extracts were

dried over sodium sulfate and concentrated under vacuum. The product was characterized by mass spectroscopy.




2-(4-aminophenyl)-7-methyl benzimidazole

2-(4-Aminophenyl)-6-methyl benzimidazole was synthesized from 2-(4-nitrophenyl)-6-methyl benzimidazole, which was prepared by mixing 1,2-diamino-4-methylbenzene (1.24 g, 9.8 mmole) with 4-nitrobenzoic acid (1.6 g, 9.9 mmole) and dissolved in POCl_3 (10 ml) and heated to reflux for 2.5 hours. The reaction mixture was cooled and cautiously poured onto ice. The resulting solid was filtered, washed with NaHCO_3 and used without further purification.



2-(4-nitrophenyl)-6-methyl benzimidazole


2-(4-Aminophenyl)-6-methyl benzimidazole was synthesized by dissolving 1 g of the above nitrobenzimidazole in 30% $\text{Na}_2\text{S} \cdot 9\text{H}_2\text{O}$ (20 ml) with stirring at RT for 4.5 h. The reaction mixture was diluted with water and extracted with EtOAc. The combined organic extracts were dried over sodium sulfate and concentrated under vacuum. The product was characterized by mass spectroscopy.



2-(4-aminophenyl)-6-methyl benzimidazole


2-(4-Aminophenyl)-5,6-dimethyl benzimidazole was synthesized from 2-(4-nitrophenyl)-5,6-dimethyl benzimidazole, which was prepared by mixing 1,2-diamino-4,5-dimethylbenzene (1.38 g, 10.1 mmole) with 4-nitrobenzoic acid (1.69 g, 9.9 mmole) and dissolved in POCl_3 (10 ml) and heated to reflux for 2.5 hours. The reaction mixture was cooled and cautiously poured

onto ice. The resulting solid was filtered, washed with NaHCO_3 and used without further purification.



2-(4-nitrophenyl)-5,6-dimethyl benzimidazole

2-(4-Aminophenyl)-5,6-dimethyl benzimidazole was synthesized by dissolving 1 g of the above nitrobenzimidazole (31.1) in 30% $\text{Na}_2\text{S}\cdot 9\text{H}_2\text{O}$ (20 ml) with stirring at RT for 4.5 h. The reaction mixture was diluted with water and extracted with EtOAc. The combined organic extracts were dried over sodium sulfate and concentrated under vacuum. The product was characterized by mass spectroscopy.



2-(4-aminophenyl)-5,6-dimethyl benzimidazole

The subsequent preparation of symmetrical diamides was accomplished by one of the following methods:

Method A: 2-(4-Aminophenyl)-6-aminobenzimidazole (1 mmole) was suspended in THF (5 ml) to which was added DIEA (2.5 mmole) and mixture cooled to -78°C . To the above cooled mixture was added the acid chloride (2.5 mmole) and let warm to RT overnight. Water (2 ml) is added to the reaction and extracted with EtOAc. The combined organic extracts were combined washed with NaHCO_3 (aq.) and concentrated under reduced pressure. The resulting residue was purified on silica gel (hexanes/EtOAc or $\text{MeOH}/\text{CH}_2\text{Cl}_2$) or reverse phase HPLC ($\text{CH}_3\text{CN}/\text{H}_2\text{O}$).

Method B: 2-(4-Aminophenyl)-6-aminobenzimidazole (1 mmole) and DMAP (cat.) was dissolved in pyridine (5 ml). To the above solution was added the acid chloride (2.5 mmole) and the reaction stirred overnight at 60°C . The reaction was cooled to room temperature and water added to precipitate the product. The resulting solid was collected by filtration with the solid

being washed by hexanes and water and NaHCO_3 (aq.). The resulting residue was purified on silica gel (hexanes/EtOAc or MeOH/ CH_2Cl_2) or reverse phase HPLC ($\text{CH}_3\text{CN}/\text{H}_2\text{O}$).

Method C: 2-(4-Aminophenyl)-6-aminobenzimidazole (1 mmole) was suspended in THF (10 ml) to which was added K_2CO_3 (2.5 mmole) in water (0.5 ml), and mixture cooled to -78°C . To the above cooled mixture was added the acid chloride (2.5 mmole) and let warm to RT overnight. Water (10 ml) was added to the reaction and extracted with EtOAc. The combined organic extracts were combined washed with NaHCO_3 (aq.) and concentrated under reduced pressure. The resulting residue was purified on silica gel (hexanes/EtOAc or MeOH/ CH_2Cl_2) or reverse phase HPLC ($\text{CH}_3\text{CN}/\text{H}_2\text{O}$).

Method D: The carboxylic acid (2.2 mmole), EDC (2.2 mmole) and DMAP (cat.) was dissolved in hot pyridine. To the above solution was added 2-(4-aminophenyl)-6-aminobenzimidazole (1 mmole) and heated to 60°C overnight. The cooled reaction mixture was partitioned between water and EtOAc. The organic layer was washed with NaHCO_3 , dried over Na_2SO_4 and concentrated under vacuum. The resulting residue was purified on silica gel (hexanes/EtOAc or MeOH/ CH_2Cl_2) or reverse phase HPLC ($\text{CH}_3\text{CN}/\text{H}_2\text{O}$).

Diacyl Benzimidazole Species

The following species encompassed within the disclosed generic formula were synthesized and tested for their ability to suppress IgE. The species are presented above in *the Summary of the Invention*

IgE Down-Regulatory Activity

All of the disclosed species were tested for their ability to suppress IgE in both the *ex vivo* and *in vivo* assays. They were all active in both assays. Activities (IC_{50}) of the species in the *ex vivo* assay ranged from about 100 pM to 1 nM. In the *in vivo* assay, the IC_{50} dose ranged from approximately 100 μ g/kg body weight/day to about 10 mg/kg body weight/day. The diacyl benzimidazole compounds were generally more potent than the monoacyl compounds.

Suppression of IgE Response

The inhibitory activity of the small molecules of the present invention were assayed using both the *ex vivo* and *in vivo* assays as described above. All of the compounds presented above were active in suppressing the IgE response. In the *ex vivo* assay, compounds in genres I-XI produced 50% inhibition at concentrations ranging from 1 pM to 10 μ M. In the *in vivo* assay, the compounds were effective at concentrations ranging from less than about 0.01 mg/kg/day to about 25 mg/kg/day, when administered in divided doses (e.g., two to four times daily) for at least two to seven consecutive days. Thus, the small molecule inhibitors of the present invention are disclosed as being useful in lowering the antigen-induced increase in IgE concentration, and consequently, in the treatment of IgE-dependent processes such as allergies in general and allergic asthma in particular.

Treatment Regimens

The amount of the IgE inhibitor compound which may be effective in treating a particular allergy or condition will depend on the nature of the disorder, and can be determined by standard clinical techniques. The precise dose to be employed in a given situation will also depend on the choice of compound and the seriousness of the condition, and should be decided according to the judgment of the practitioner and each patient's circumstances. Appropriate dosages can be determined and adjusted by the practitioner based on dose response relationships between the patient's IgE levels as well as standard indices of pulmonary and hemodynamic changes. Moreover, those skilled in the art will appreciate that dose ranges can be determined without undue experimentation by following the protocol(s) disclosed herein for *ex vivo* and *in vivo* screening (See

for example Hasegawa et al., *J. Med. Chem.* 40: 395-407 (1997) and Ohmori et al., *Int. J. Immunopharmacol.* 15:573-579 (1993); employing similar *ex vivo* and *in vivo* assays for determining dose-response relationships for IgE suppression by naphthalene derivatives; incorporated herein by reference).


Initially, suitable dosages of the compounds will generally range from about 0.001 mg to about 300 mg per kg body weight per day in divided doses, more preferably, between about 0.01 mg and 100 mg per kg body weight per day in divided doses. The compounds are preferably administered systemically as pharmaceutical formulations appropriate to such routes as oral, aerosol, intravenous, subcutaneously, or by any other route which may be effective in providing systemic dosing of the active compound. The compositions of pharmaceutical formulations are well known in the art. The treatment regimen preferably involves periodic administration. Moreover, long-term therapy may be indicated where allergic reactions appear to be triggered by continuous exposure to the allergen(s). Daily or twice daily administration has been effective in suppressing the IgE response to a single antigen challenge in animals when carried out continuously from a period of two to seven consecutive days. Thus, in a preferred embodiment, the compound is administered for at least two consecutive days at regular periodic intervals. However, the treatment regimen, including frequency of dosing and duration of treatment may be determined by the skilled practitioner, and modified as needed to provide optimal IgE down-regulation, depending on nature of the allergen, the dose, frequency, and duration of the allergen exposure, and the standard clinical indices.

In one embodiment of the present invention, an IgE-suppressing compound may be administered in conjunction with one or more of the other small molecule inhibitors disclosed, in order to produce optimal down-regulation of the patient's IgE response. Further, it is envisioned that one or more of the compounds of the present invention may be administered in combination with other drugs already known or later discovered for treatment of the underlying cause as well as the acute symptoms of allergy or asthma. Such combination therapies envisioned within the scope of the present invention include mixing of one or more of the small molecule IgE-inhibitors together with one or more additional ingredients, known to be effective in reducing at least one symptom of the disease condition. In a variation, the small molecule IgE-inhibitors herein disclosed may be administered separately from the additional drugs, but during the same course of the disease

condition, wherein both the IgE-inhibitor(s) and the palliative compounds are administered in accordance with their independent effective treatment regimens.

WHAT IS CLAIMED IS:

1. A pharmaceutical composition comprising the following compounds:















wherein X and Y are independently selected from the group consisting of H, alkyl, alkoxy, aryl, substituted aryl, hydroxy, halogen, amino, alkylamino, nitro, cyano, CF₃, OCF₃, CONH₂, CONHR and NHCOR₁;













wherein R is selected from the group consisting of H, CH₃, C₂H₅, C₃H₇, C₄H₉, CH₂Ph, and CH₂C₆H₄-F(p-); and













wherein R₁ and R₂ are independently selected from the group consisting of H, aryl, substituted aryl; cycloaryl substituted cycloaryl, multi-ring cycloaryl, benzyl, substituted benzyl, alkyl, cycloalkyl substituted cycloalkyl, multi-ring cycloalkyl, fused-ring aliphatic, cyclopropyl; substituted cyclopropyl, cyclobutyl, substituted cyclobutyl, cyclopentyl, substituted cyclopentyl, cyclohexyl, substituted cyclohexyl, cycloheptyl, substituted cycloheptyl, bicycloheptyl, bicyclooctyl, bicyclononyl, substituted bicycloalkenyl, adamantyl, substituted adamantyl and the like, wherein at least one of R₁ and R₂ are aromatic groups.













2. The pharmaceutical composition of claim 1, wherein the R₁ and R₂ substitutions are selected from the group consisting of alkyl, aryl, CF₃, CH₃, OCH₃, OH, CN, COOR and COOH.













3. The pharmaceutical composition of Claim 2, wherein the compound is selected from the group consisting of:









































































Structure 	BIGA
	$C_{27}H_{18}Cl_2N_4O_2$
	CLOGP 5.34


Structure 	BIGB
	$C_{27}H_{17}Cl_3N_4O_2$
	CLOGP 6.12


Structure 	BIGC
	$C_{28}H_{20}Cl_2N_4O_3$
	CLOGP 5.46


Structure 	BIGD
	$C_{28}H_{17}Cl_2N_4O_2$
	CLOGP 4.16


Structure 	BIGE
	$C_{30}H_{24}Cl_2N_4O_5$
	CLOGP 4.69


Structure 	BIGF
	$C_{27}H_{16}Cl_4N_4O_2$
	CLOGP 6.85


Structure 	BIGG
	$C_{27}H_{16}Cl_4N_4O_2$
	CLOGP 5.19













Structure 	BIGH
	$C_{27}H_{16}Cl_4N_4O_2$
	CLOGP 6.02













Structure 	BIGI
	$C_{27}H_{24}Cl_2N_4O_2$
	CLOGP 5.88













Structure 	BIGJ
	$C_{26}H_{22}Cl_2N_4O_2$
	CLOGP 5.33













Structure 	BIGK
	$C_{25}H_{16}Cl_2N_4O_2$ S
	CLOGP 5.08













Structure 	BIGL
	$C_{25}H_{16}Cl_2N_4O_2$ S
	CLOGP 5.08

















Structure 	BIJG
	$C_{26}H_{22}Cl_2N_4O_2$
	CLOGP 5.29


Structure 	BIJH
	$C_{26}H_{22}Cl_2N_4O_2$
	CLOGP 6.12


Structure 	BIIA
	$C_{27}H_{26}N_4O_2$
	CLOGP 6.01


Structure 	BIIB
	$C_{27}H_{25}ClN_4O_2$
	CLOGP 6.78


Structure 	BIIC
	$C_{28}H_{28}N_4O_3$
	CLOGP 6.12


Structure 	BIID
	$C_{26}H_{25}N_5O_2$
	CLOGP 4.82


Structure 	BII E
	$C_{30}H_{32}N_4O_5$
	CLOGP 5.36


Structure 	BIIF
	$C_{27}H_{24}Cl_2N_4O_2$
	CLOGP 7.51


Structure 	BIIG
	$C_{27}H_{24}Cl_2N_4O_2$
	CLOGP 5.85


Structure 	BIIH
	$C_{27}H_{24}Cl_2N_4O_2$
	CLOGP 6.68


Structure 	BIJK
	$C_{25}H_{24}N_4O_2S$
	CLOGP 5.74


Structure 	BIIL
	$C_{25}H_{24}N_4O_2S$
	CLOGP 5.74


Structure 	BIJA
	$C_{26}H_{24}N_4O_2$
	CLOGP 5.45













Structure 	BIJB
	$C_{26}H_{23}ClN_4O_2$
	CLOGP 6.22











Structure 	BIJC
	$C_{27}H_{26}N_4O_3$
	CLOGP 5.56



Structure 	BIJD
	$C_{25}H_{23}N_5O_2$
	CLOGP 4.26

Structure 	BIJE
	$C_{29}H_{30}N_4O_5$
	CLOGP 4.80









Structure 	BIJF
	$C_{26}H_{22}Cl_2N_4O_2$
	CLOGP 6.95











avanir_vlib.db





Structure 	BIPG
	$C_{28}H_{26}Cl_2N_4O_2$
	CLOGP 6.41


Structure 	BIPH
	$C_{28}H_{26}Cl_2N_4O_2$
	CLOGP 7.24


Structure 	BIPK
	$C_{26}H_{26}N_4O_2S$
	CLOGP 6.30


Structure 	BIPL
	$C_{26}H_{26}N_4O_2S$
	CLOGP 6.30


Structure 	BIPA
	$C_{28}H_{28}N_4O_2$
	CLOGP 6.57











Structure 	BIPB
	$C_{28}H_{27}ClN_4O_2$
	CLOGP 7.34











Structure 	BIPC
	$C_{29}H_{30}N_4O_3$
	CLOGP 6.68

Structure 	BIPD
	$C_{27}H_{27}N_5O_2$
	CLOGP 5.38

Structure 	BIPE
	$C_{31}H_{34}N_4O_5$
	CLOGP 5.92

Structure 	BIPF
	$C_{28}H_{26}Cl_2N_4O_2$
	CLOGP 8.07





4. The pharmaceutical composition of any of Claims 1-3 for use in the treatment of a disease condition associated with excess IgE.

5. The pharmaceutical composition of Claim 4, further comprising at least one additional ingredient which is active in reducing at least one symptom associated with the disease condition associated with excess IgE.

6. The pharmaceutical composition of Claim 5, wherein said at least one additional ingredient is selected from the group consisting of a short-acting β_2 -adrenergic agonist, a long-acting β_2 -adrenergic agonist, an antihistamine, a phosphodiesterase inhibitor, an anticholinergic agent, a corticosteroid, an inflammatory mediator release inhibitor and a leukotriene receptor antagonist.

7. Use of the pharmaceutical composition of any one of Claims 1-3 in the preparation of a medicament for treatment of a disease condition associated with excess IgE.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/11322

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A61K31/415

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 719 765 A (MITSUI TOATSU CHEMICALS) 3 July 1996 (1996-07-03) page 30 page 31 page 38 page 39 page 49 page 50; claim 1; examples 43,88,1100,2100	1-4

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

1 October 1999

Date of mailing of the international search report

11/10/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Orviz Diaz, P

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 99/11322

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
See FURTHER INFORMATION SHEET PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International Application No. PCT/US 99/11322

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

The substituents in the general formula of claim 1 are not clearly defined, contrary to Art. 6 PCT. The expressions "the like" or "substituted aryl", for example, encompass an extremely large number of possibilities, which makes impossible to carry out a complete search.

Furthermore, most of the specific R1 and R2 substituents mentioned in claim 2 are not covered by claim 1 and some of the compounds mentioned in claim 3 have a pyridine ring or a thiophene ring, which are not mentioned as possible substituents in claim 1.

In view of this the search had to be limited to the general structural characteristics of the formula in claim 1.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/11322

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0719765 A	03-07-1996	JP 8231514 A	10-09-1996
		US 5821258 A	13-10-1998